



# Survey

## Complications associated with infant male circumcision

Infant male circumcision is performed for personal, religious, medical or cultural reasons. Progress has been achieved in the techniques used and pain control offered to infants. Published articles present arguments supporting and opposing neonatal circumcision. Complications of this procedure still occur. This survey will evaluate if CPSP participants are seeing complications arising from infant male circumcision.

**Your contribution is greatly appreciated.**

1. I am: a paediatrician \_\_\_ a paediatric subspecialist (specify) \_\_\_\_\_  
 other (specify) \_\_\_\_\_

2. Over the past 12 months, have you seen any infants with circumcision-related complications?  
 Yes \_\_\_ No \_\_\_ **If no, thank you for completing this survey.**

**If yes, check all that apply (photocopies should be used and attached to this survey if reporting more than one case per complication).**

Complications	# of cases	Resulted in hospitalization		Technique used if known (i.e., Gomco, Plastibell, Mogen)	Type of clinician who performed circumcision if known (i.e., family physician, paediatrician, surgeon, non-medical/cultural official)
		Yes	No		
Local infections requiring antibiotics		___	___		
Herpes simplex virus infection		___	___		
Degloving (excessive removal of skin)		___	___		
Bleeding requiring transfusion		___	___		
Urinary tract infection		___	___		
Septicemia		___	___		
Amputation		___	___		
Permanent disability, specify: _____		___	___		
Death		___	___		
Other, specify: _____		___	___		

3. Did you refer any patients to a surgical specialist for complications of circumcision? Yes \_\_\_ No \_\_\_  
 If yes, how many? \_\_\_\_\_

4. How many parents initiated their own concerns about their child's circumcision? \_\_\_\_\_

**Please return this survey with your monthly reporting form.  
 Thank you for your participation.**